

Automorphism groups and Ramsey properties of sparse graphs.

David Evans
Dept. of Mathematics, Imperial College London.

Joint work with Jan Hubička and Jaroslav Nešetřil

THEMES:

- Automorphism groups of nice model-theoretic structures acting on compact Hausdorff spaces.
- Connection with structural Ramsey theory (Kechris - Pestov - Todorčević Correspondence)
- Sparse graphs constructed using Hrushovski amalgamations exhibit interesting new phenomena.

THEOREM A: There is a countable ω -categorical structure M with the property that if $H \leq \text{Aut}(M)$ is (extremely) amenable, then H has infinitely many orbits on M^2 .

NOTE: By the Ryll-Nardzewski Theorem, $\text{Aut}(M)$ has finitely many orbits on M^n for all $n \in \mathbb{N}$.

Joint work with Jan Hubička and Jaroslav Nešetřil

THEMES:

- Automorphism groups of nice model-theoretic structures acting on compact Hausdorff spaces.
- Connection with structural Ramsey theory (Kechris - Pestov - Todorčević Correspondence)
- Sparse graphs constructed using Hrushovski amalgamations exhibit interesting new phenomena.

THEOREM A: There is a countable ω -categorical structure M with the property that if $H \leq \text{Aut}(M)$ is (extremely) amenable, then H has infinitely many orbits on M^2 .

NOTE: By the Ryll-Nardzewski Theorem, $\text{Aut}(M)$ has finitely many orbits on M^n for all $n \in \mathbb{N}$.

Amalgamation classes and Fraïssé limits.

L a 1st-order relational language and M a countable L -structure.

$\text{Age}(M)$: class of isomorphism types of finite substructures.

M is *homogeneous* if all isomorphism between finite substructures of M extend to automorphisms of M . In this case $\mathcal{C} = \text{Age}(M)$ satisfies:

AMALGAMATION PROPERTY (AP): If $f_1 : A \rightarrow B_1$ and $f_2 : A \rightarrow B_2$ are embeddings between elements of \mathcal{C} , then there is $C \in \mathcal{C}$ and embeddings $g_i : B_i \rightarrow C$ with $g_1 \circ f_1 = g_2 \circ f_2$.

Conversely: if \mathcal{C} is a countable class of isomorphism types of finite L -structures which is closed under taking substructures, has the joint embedding property and

\mathcal{C} has AP,

then there is a countable, homogeneous structure $M(\mathcal{C})$ with $\text{Age}(M(\mathcal{C})) = \mathcal{C}$. It is unique up to isomorphism.

\mathcal{C} is an *amalgamation class* and $M(\mathcal{C})$ is its *Fraïssé limit*.

Amalgamation classes and Fraïssé limits.

L a 1st-order relational language and M a countable L -structure.

$\text{Age}(M)$: class of isomorphism types of finite substructures.

M is *homogeneous* if all isomorphism between finite substructures of M extend to automorphisms of M . In this case $\mathcal{C} = \text{Age}(M)$ satisfies:

AMALGAMATION PROPERTY (AP): If $f_1 : A \rightarrow B_1$ and $f_2 : A \rightarrow B_2$ are embeddings between elements of \mathcal{C} , then there is $C \in \mathcal{C}$ and embeddings $g_j : B_j \rightarrow C$ with $g_1 \circ f_1 = g_2 \circ f_2$.

Conversely: if \mathcal{C} is a countable class of isomorphism types of finite L -structures which is closed under taking substructures, has the joint embedding property and

\mathcal{C} has AP,

then there is a countable, homogeneous structure $M(\mathcal{C})$ with $\text{Age}(M(\mathcal{C})) = \mathcal{C}$. It is unique up to isomorphism.

\mathcal{C} is an *amalgamation class* and $M(\mathcal{C})$ is its *Fraïssé limit*.

Amalgamation classes and Fraïssé limits.

L a 1st-order relational language and M a countable L -structure.

$\text{Age}(M)$: class of isomorphism types of finite substructures.

M is *homogeneous* if all isomorphism between finite substructures of M extend to automorphisms of M . In this case $\mathcal{C} = \text{Age}(M)$ satisfies:

AMALGAMATION PROPERTY (AP): If $f_1 : A \rightarrow B_1$ and $f_2 : A \rightarrow B_2$ are embeddings between elements of \mathcal{C} , then there is $C \in \mathcal{C}$ and embeddings $g_i : B_i \rightarrow C$ with $g_1 \circ f_1 = g_2 \circ f_2$.

Conversely: if \mathcal{C} is a countable class of isomorphism types of finite L -structures which is closed under taking substructures, has the joint embedding property and

\mathcal{C} has AP,

then there is a countable, homogeneous structure $M(\mathcal{C})$ with $\text{Age}(M(\mathcal{C})) = \mathcal{C}$. It is unique up to isomorphism.

\mathcal{C} is an *amalgamation class* and $M(\mathcal{C})$ is its *Fraïssé limit*.

EXAMPLE:

\mathcal{G} the class of all finite graphs; $M(\mathcal{G})$ is the Random Graph.

VARIATION: Can also work with a distinguished notion of embedding / substructure, $(\mathcal{C}; \leq)$.

– This is used in the Hrushovski construction.

EXAMPLE:

\mathcal{G} the class of all finite graphs; $M(\mathcal{G})$ is the Random Graph.

VARIATION: Can also work with a distinguished notion of embedding / substructure, $(\mathcal{C}; \leq)$.

– This is used in the Hrushovski construction.

Ramsey classes

L^{\leq} : relational language with \leq .

\mathcal{A} : a class of finite L^{\leq} -structures closed under substrs and satisfying JEP and where \leq is a linear ordering.

DEFINITION: Say that \mathcal{A} is a **Ramsey class** if whenever $A \subseteq B \in \mathcal{A}$, there is $B \subseteq C \in \mathcal{A}$ such that if

$$\gamma: \binom{C}{A} \rightarrow \{0, 1\}$$

is a 2-colouring of the copies of A in C , there is $B' \in \binom{C}{B}$ (a copy of B in C) such that γ is constant on $\binom{B'}{A}$.

EXAMPLES: (1) $L = \{\leq\}$. Take $\mathcal{A} =$ finite linear orders.

(2) (Nešetřil - Rödl) The class \mathcal{G}^{\leq} of linearly ordered finite graphs.

THEOREM: (Nešetřil) If \mathcal{A} is a Ramsey class, then \mathcal{A} has the amalgamation property.

– What's special about $M(\mathcal{A})$?

Ramsey classes

L^{\leq} : relational language with \leq .

\mathcal{A} : a class of finite L^{\leq} -structures closed under substrs and satisfying JEP and where \leq is a linear ordering.

DEFINITION: Say that \mathcal{A} is a **Ramsey class** if whenever $A \subseteq B \in \mathcal{A}$, there is $B \subseteq C \in \mathcal{A}$ such that if

$$\gamma: \binom{C}{A} \rightarrow \{0, 1\}$$

is a 2-colouring of the copies of A in C , there is $B' \in \binom{C}{B}$ (a copy of B in C) such that γ is constant on $\binom{B'}{A}$.

EXAMPLES: (1) $L = \{\leq\}$. Take $\mathcal{A} =$ finite linear orders.

(2) (Nešetřil - Rödl) The class \mathcal{G}^{\leq} of linearly ordered finite graphs.

THEOREM: (Nešetřil) If \mathcal{A} is a Ramsey class, then \mathcal{A} has the amalgamation property.

– What's special about $M(\mathcal{A})$?

Ramsey classes

L^{\leq} : relational language with \leq .

\mathcal{A} : a class of finite L^{\leq} -structures closed under substrs and satisfying JEP and where \leq is a linear ordering.

DEFINITION: Say that \mathcal{A} is a **Ramsey class** if whenever $A \subseteq B \in \mathcal{A}$, there is $B \subseteq C \in \mathcal{A}$ such that if

$$\gamma: \binom{C}{A} \rightarrow \{0, 1\}$$

is a 2-colouring of the copies of A in C , there is $B' \in \binom{C}{B}$ (a copy of B in C) such that γ is constant on $\binom{B'}{A}$.

EXAMPLES: (1) $L = \{\leq\}$. Take $\mathcal{A} =$ finite linear orders.

(2) (Nešetřil - Rödl) The class \mathcal{G}^{\leq} of linearly ordered finite graphs.

THEOREM: (Nešetřil) If \mathcal{A} is a Ramsey class, then \mathcal{A} has the amalgamation property.

– What's special about $M(\mathcal{A})$?

Ramsey classes

L^{\leq} : relational language with \leq .

\mathcal{A} : a class of finite L^{\leq} -structures closed under substrs and satisfying JEP and where \leq is a linear ordering.

DEFINITION: Say that \mathcal{A} is a **Ramsey class** if whenever $A \subseteq B \in \mathcal{A}$, there is $B \subseteq C \in \mathcal{A}$ such that if

$$\gamma: \binom{C}{A} \rightarrow \{0, 1\}$$

is a 2-colouring of the copies of A in C , there is $B' \in \binom{C}{B}$ (a copy of B in C) such that γ is constant on $\binom{B'}{A}$.

EXAMPLES: (1) $L = \{\leq\}$. Take $\mathcal{A} =$ finite linear orders.

(2) (Nešetřil - Rödl) The class \mathcal{G}^{\leq} of linearly ordered finite graphs.

THEOREM: (Nešetřil) If \mathcal{A} is a Ramsey class, then \mathcal{A} has the amalgamation property.

– What's special about $M(\mathcal{A})$?

Ramsey classes

L^{\leq} : relational language with \leq .

\mathcal{A} : a class of finite L^{\leq} -structures closed under substrs and satisfying JEP and where \leq is a linear ordering.

DEFINITION: Say that \mathcal{A} is a **Ramsey class** if whenever $A \subseteq B \in \mathcal{A}$, there is $B \subseteq C \in \mathcal{A}$ such that if

$$\gamma: \binom{C}{A} \rightarrow \{0, 1\}$$

is a 2-colouring of the copies of A in C , there is $B' \in \binom{C}{B}$ (a copy of B in C) such that γ is constant on $\binom{B'}{A}$.

EXAMPLES: (1) $L = \{\leq\}$. Take $\mathcal{A} =$ finite linear orders.

(2) (Nešetřil - Rödl) The class \mathcal{G}^{\leq} of linearly ordered finite graphs.

THEOREM: (Nešetřil) If \mathcal{A} is a Ramsey class, then \mathcal{A} has the amalgamation property.

– What's special about $M(\mathcal{A})$?

Automorphism groups.

Ω infinite set (usually countable); $\text{Sym}(\Omega)$ symmetric group.

$G \leq \text{Sym}(\Omega) \subseteq \Omega^\Omega$ pointwise convergence topology.

Basic open sets: $\{g \in G : g|A = \gamma\}$, $A \subseteq \Omega$ finite and $\gamma : A \rightarrow \Omega$.

G is a topological group.

$\text{Sym}(\Omega)$ complete metrizable if Ω is countable.

Lemma

$G \leq \text{Sym}(\Omega)$ is closed iff $G = \text{Aut}(M)$ for some 1st order structure M with domain Ω .

INTERESTING EXAMPLES: M countable homogeneous, or ω -categorical.

REMARK: If $G \leq \text{Sym}(\Omega)$ is closed there is a *homogeneous* structure M with $\text{Aut}(M) = G$ (but the language may have to be infinite).

Automorphism groups.

Ω infinite set (usually countable); $\text{Sym}(\Omega)$ symmetric group.

$G \leq \text{Sym}(\Omega) \subseteq \Omega^\Omega$ pointwise convergence topology.

Basic open sets: $\{g \in G : g|A = \gamma\}$, $A \subseteq \Omega$ finite and $\gamma : A \rightarrow \Omega$.

G is a topological group.

$\text{Sym}(\Omega)$ complete metrizable if Ω is countable.

Lemma

$G \leq \text{Sym}(\Omega)$ is closed iff $G = \text{Aut}(M)$ for some 1st order structure M with domain Ω .

INTERESTING EXAMPLES: M countable homogeneous, or ω -categorical.

REMARK: If $G \leq \text{Sym}(\Omega)$ is closed there is a *homogeneous* structure M with $\text{Aut}(M) = G$ (but the language may have to be infinite).

Automorphism groups.

Ω infinite set (usually countable); $\text{Sym}(\Omega)$ symmetric group.

$G \leq \text{Sym}(\Omega) \subseteq \Omega^\Omega$ pointwise convergence topology.

Basic open sets: $\{g \in G : g|A = \gamma\}$, $A \subseteq \Omega$ finite and $\gamma : A \rightarrow \Omega$.

G is a topological group.

$\text{Sym}(\Omega)$ complete metrizable if Ω is countable.

Lemma

$G \leq \text{Sym}(\Omega)$ is closed iff $G = \text{Aut}(M)$ for some 1st order structure M with domain Ω .

INTERESTING EXAMPLES: M countable homogeneous, or ω -categorical.

REMARK: If $G \leq \text{Sym}(\Omega)$ is closed there is a *homogeneous* structure M with $\text{Aut}(M) = G$ (but the language may have to be infinite).

Automorphism groups.

Ω infinite set (usually countable); $\text{Sym}(\Omega)$ symmetric group.

$G \leq \text{Sym}(\Omega) \subseteq \Omega^\Omega$ pointwise convergence topology.

Basic open sets: $\{g \in G : g|A = \gamma\}$, $A \subseteq \Omega$ finite and $\gamma : A \rightarrow \Omega$.

G is a topological group.

$\text{Sym}(\Omega)$ complete metrizable if Ω is countable.

Lemma

$G \leq \text{Sym}(\Omega)$ is closed iff $G = \text{Aut}(M)$ for some 1st order structure M with domain Ω .

INTERESTING EXAMPLES: M countable homogeneous, or ω -categorical.

REMARK: If $G \leq \text{Sym}(\Omega)$ is closed there is a *homogeneous* structure M with $\text{Aut}(M) = G$ (but the language may have to be infinite).

Automorphism groups.

Ω infinite set (usually countable); $\text{Sym}(\Omega)$ symmetric group.

$G \leq \text{Sym}(\Omega) \subseteq \Omega^\Omega$ pointwise convergence topology.

Basic open sets: $\{g \in G : g|A = \gamma\}$, $A \subseteq \Omega$ finite and $\gamma : A \rightarrow \Omega$.

G is a topological group.

$\text{Sym}(\Omega)$ complete metrizable if Ω is countable.

Lemma

$G \leq \text{Sym}(\Omega)$ is closed iff $G = \text{Aut}(M)$ for some 1st order structure M with domain Ω .

INTERESTING EXAMPLES: M countable homogeneous, or ω -categorical.

REMARK: If $G \leq \text{Sym}(\Omega)$ is closed there is a *homogeneous* structure M with $\text{Aut}(M) = G$ (but the language may have to be infinite).

Topological Dynamics

G a topological group.

G -flow: compact, Hausdorff, non-empty space X with a continuous G -action.

Definition

- 1 G is *amenable* if every G -flow X supports a G -invariant Borel probability measure.
- 2 G is *extremely amenable* if every G -flow has a fixed point.

Topological Dynamics

G a topological group.

G -flow: compact, Hausdorff, non-empty space X with a continuous G -action.

Definition

- 1 G is *amenable* if every G -flow X supports a G -invariant Borel probability measure.
- 2 G is *extremely amenable* if every G -flow has a fixed point.

Topological Dynamics

G a topological group.

G -flow: compact, Hausdorff, non-empty space X with a continuous G -action.

Definition

- 1 G is *amenable* if every G -flow X supports a G -invariant Borel probability measure.
- 2 G is *extremely amenable* if every G -flow has a fixed point.

Topological Dynamics

G a topological group.

G -flow: compact, Hausdorff, non-empty space X with a continuous G -action.

Definition

- 1 G is *amenable* if every G -flow X supports a G -invariant Borel probability measure.
- 2 G is *extremely amenable* if every G -flow has a fixed point.

G-flows

$G = \text{Aut}(M)$. Some G -flows:

- 1 Take G -invariant $\Delta \subseteq M^n$; consider $Y = \{0, 1\}^\Delta$ as a G -flow. Also consider G -invariant, closed subspaces X of Y .
- 2 G -invariant, closed subspaces of $S(M)$, Stone space over M .

EXAMPLE: $G = \text{Sym}(\Omega)$. We have a G -flow:

$$LO(\Omega) = \{R \subseteq \Omega^2 : R \text{ is a linear order on } \Omega\}.$$

COROLLARY: If $H \leq G$ is e.a. then there is an H -invariant linear order on Ω .

Theorem (Pestov, 1998)

$\text{Aut}(\mathbb{Q}; \leq)$ is e.a.

G-flows

$G = \text{Aut}(M)$. Some G -flows:

- 1 Take G -invariant $\Delta \subseteq M^n$; consider $Y = \{0, 1\}^\Delta$ as a G -flow. Also consider G -invariant, closed subspaces X of Y .
- 2 G -invariant, closed subspaces of $S(M)$, Stone space over M .

EXAMPLE: $G = \text{Sym}(\Omega)$. We have a G -flow:

$$LO(\Omega) = \{R \subseteq \Omega^2 : R \text{ is a linear order on } \Omega\}.$$

COROLLARY: If $H \leq G$ is e.a. then there is an H -invariant linear order on Ω .

Theorem (Pestov, 1998)

$\text{Aut}(\mathbb{Q}; \leq)$ is e.a.

G-flows

$G = \text{Aut}(M)$. Some G -flows:

- 1 Take G -invariant $\Delta \subseteq M^n$; consider $Y = \{0, 1\}^\Delta$ as a G -flow. Also consider G -invariant, closed subspaces X of Y .
- 2 G -invariant, closed subspaces of $S(M)$, Stone space over M .

EXAMPLE: $G = \text{Sym}(\Omega)$. We have a G -flow:

$$LO(\Omega) = \{R \subseteq \Omega^2 : R \text{ is a linear order on } \Omega\}.$$

COROLLARY: If $H \leq G$ is e.a. then there is an H -invariant linear order on Ω .

Theorem (Pestov, 1998)

$\text{Aut}(\mathbb{Q}; \leq)$ is e.a.

G-flows

$G = \text{Aut}(M)$. Some G -flows:

- 1 Take G -invariant $\Delta \subseteq M^n$; consider $Y = \{0, 1\}^\Delta$ as a G -flow. Also consider G -invariant, closed subspaces X of Y .
- 2 G -invariant, closed subspaces of $S(M)$, Stone space over M .

EXAMPLE: $G = \text{Sym}(\Omega)$. We have a G -flow:

$$LO(\Omega) = \{R \subseteq \Omega^2 : R \text{ is a linear order on } \Omega\}.$$

COROLLARY: If $H \leq G$ is e.a. then there is an H -invariant linear order on Ω .

Theorem (Pestov, 1998)

$\text{Aut}(\mathbb{Q}; \leq)$ is e.a.

G-flows

$G = \text{Aut}(M)$. Some G -flows:

- 1 Take G -invariant $\Delta \subseteq M^n$; consider $Y = \{0, 1\}^\Delta$ as a G -flow. Also consider G -invariant, closed subspaces X of Y .
- 2 G -invariant, closed subspaces of $S(M)$, Stone space over M .

EXAMPLE: $G = \text{Sym}(\Omega)$. We have a G -flow:

$$LO(\Omega) = \{R \subseteq \Omega^2 : R \text{ is a linear order on } \Omega\}.$$

COROLLARY: If $H \leq G$ is e.a. then there is an H -invariant linear order on Ω .

Theorem (Pestov, 1998)

$\text{Aut}(\mathbb{Q}; \leq)$ is e.a.

The Kechris - Pestov - Todorčević Correspondence

Theorem (KPT, 2005)

Suppose M is a countable, homogeneous, linearly ordered relational structures with age \mathcal{A} . TFAE:

- 1 $\text{Aut}(M)$ is extremely amenable.
- 2 \mathcal{A} is a Ramsey class.

So Ramsey classes correspond to homogeneous structures with e.a. automorphism groups.

EXAMPLE: \mathcal{G}^{\leq} (finite l.o. graphs) is a Ramsey class. Let $\Gamma^{\leq} = M(\mathcal{G}^{\leq})$. Then $\text{Aut}(\Gamma^{\leq})$ is e.a. The graph reduct Γ is the Random Graph and $\text{Aut}(\Gamma^{\leq}) \leq \text{Aut}(\Gamma)$.

Note that \mathcal{G}^{\leq} is a precompact expansion of \mathcal{G} : every $A \in \mathcal{G}$ expands to finitely many iso types of structures in \mathcal{G}^{\leq} .

Equivalently each $\text{Aut}(\Gamma)$ -orbit on Γ^n splits into finitely many $\text{Aut}(\Gamma^{\leq})$ -orbits.

The universal minimal flow

A G -flow X is *minimal* if every G -orbit on X is dense.

FACT: (Ellis) There is a unique *universal* minimal G -flow, $M(G)$.

DEF: Let $G = \text{Aut}(M)$. Say $H \leq G$ is *precompact* if for every G -orbit $\Delta \subseteq M^n$, H has finitely many orbits on Δ .

KPT; Nguyen Van Thé

Suppose M is a countable L -structure. If $G = \text{Aut}(M)$ has a precompact e.a. closed subgroup $H = \text{Aut}(N)$, then $M(G)$ can be described. In particular, $M(G)$ is metrizable and has a comeagre orbit. The same is therefore true of every minimal G -flow.

EXAMPLES: (1) $M(\text{Sym}(\Omega)) = LO(\Omega)$.

(2) If Γ is the random graph, then $M(\text{Aut}(\Gamma)) = LO(\Gamma)$.

COROLLARY: $\text{Sym}(\Omega)$ and $\text{Aut}(\Gamma)$ are amenable.

The universal minimal flow

A G -flow X is *minimal* if every G -orbit on X is dense.

FACT: (Ellis) There is a unique *universal* minimal G -flow, $M(G)$.

DEF: Let $G = \text{Aut}(M)$. Say $H \leq G$ is *precompact* if for every G -orbit $\Delta \subseteq M^n$, H has finitely many orbits on Δ .

KPT; Nguyen Van Thé

Suppose M is a countable L -structure. If $G = \text{Aut}(M)$ has a precompact e.a. closed subgroup $H = \text{Aut}(N)$, then $M(G)$ can be described. In particular, $M(G)$ is metrizable and has a comeagre orbit. The same is therefore true of every minimal G -flow.

EXAMPLES: (1) $M(\text{Sym}(\Omega)) = LO(\Omega)$.

(2) If Γ is the random graph, then $M(\text{Aut}(\Gamma)) = LO(\Gamma)$.

COROLLARY: $\text{Sym}(\Omega)$ and $\text{Aut}(\Gamma)$ are amenable.

The universal minimal flow

A G -flow X is *minimal* if every G -orbit on X is dense.

FACT: (Ellis) There is a unique *universal* minimal G -flow, $M(G)$.

DEF: Let $G = \text{Aut}(M)$. Say $H \leq G$ is *precompact* if for every G -orbit $\Delta \subseteq M^n$, H has finitely many orbits on Δ .

KPT; Nguyen Van Thé

Suppose M is a countable L -structure. If $G = \text{Aut}(M)$ has a precompact e.a. closed subgroup $H = \text{Aut}(N)$, then $M(G)$ can be described. In particular, $M(G)$ is metrizable and has a comeagre orbit. The same is therefore true of every minimal G -flow.

EXAMPLES: (1) $M(\text{Sym}(\Omega)) = LO(\Omega)$.

(2) If Γ is the random graph, then $M(\text{Aut}(\Gamma)) = LO(\Gamma)$.

COROLLARY: $\text{Sym}(\Omega)$ and $\text{Aut}(\Gamma)$ are amenable.

The universal minimal flow

A G -flow X is *minimal* if every G -orbit on X is dense.

FACT: (Ellis) There is a unique *universal* minimal G -flow, $M(G)$.

DEF: Let $G = \text{Aut}(M)$. Say $H \leq G$ is *precompact* if for every G -orbit $\Delta \subseteq M^n$, H has finitely many orbits on Δ .

KPT; Nguyen Van Thé

Suppose M is a countable L -structure. If $G = \text{Aut}(M)$ has a precompact e.a. closed subgroup $H = \text{Aut}(N)$, then $M(G)$ can be described. In particular, $M(G)$ is metrizable and has a comeagre orbit. The same is therefore true of every minimal G -flow.

EXAMPLES: (1) $M(\text{Sym}(\Omega)) = LO(\Omega)$.

(2) If Γ is the random graph, then $M(\text{Aut}(\Gamma)) = LO(\Gamma)$.

COROLLARY: $\text{Sym}(\Omega)$ and $\text{Aut}(\Gamma)$ are amenable.

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Question

- Question asked (around 2011) by: Bodirsky, Pinsker, Tsankov; Nešetřil; Nguyen van Thé:
 - ▶ If M is countable ω -categorical, is there an ω -categorical expansion N of M with $\text{Aut}(N)$ extremely amenable? Equivalently, is there a precompact e.a. closed subgroup of $\text{Aut}(M)$.
- Particularly interesting case: M homogeneous in a finite relational language.
- Why ask the question?
 - ▶ Ubiquity of ω -categorical structures with e.a. automorphism groups
 - ▶ Ubiquity of Ramsey classes
 - ▶ Applications: reducts; complexity of CSP's (Bodirsky, Pinsker et al.)
 - ▶ Describing $M(G)$ for G closed, oligomorphic permutation group.
 - ▶ Evidence. Work on Ramsey expansions of Fraïssé classes: Nešetřil - Rödl; Jasinski, Laflamme, Nguyen van Thé, Woodrow; ...

Sparse graphs.

DEF: Suppose $k \in \mathbb{N}$. A graph $M = (M; E)$ is k -sparse if for all finite $A \subseteq M$ we have $|E[A]| \leq k|A|$.

FACT: If the graph $M = (M; E)$ is k -sparse, then it is k -orientable: the edges of M can be directed so that each vertex has at most k directed edges coming out.

DEF: If M is k -sparse, let

$$X(M) = \{D \subseteq M^2 : (M; D) \text{ is a } k\text{-orientation of } M\} \subseteq \{0, 1\}^{M^2}.$$

Note that this is an $\text{Aut}(M)$ -flow.

Sparse graphs.

DEF: Suppose $k \in \mathbb{N}$. A graph $M = (M; E)$ is k -sparse if for all finite $A \subseteq M$ we have $|E[A]| \leq k|A|$.

FACT: If the graph $M = (M; E)$ is k -sparse, then it is k -orientable: the edges of M can be directed so that each vertex has at most k directed edges coming out.

DEF: If M is k -sparse, let

$$X(M) = \{D \subseteq M^2 : (M; D) \text{ is a } k\text{-orientation of } M\} \subseteq \{0, 1\}^{M^2}.$$

Note that this is an $\text{Aut}(M)$ -flow.

Sparse graphs.

DEF: Suppose $k \in \mathbb{N}$. A graph $M = (M; E)$ is k -sparse if for all finite $A \subseteq M$ we have $|E[A]| \leq k|A|$.

FACT: If the graph $M = (M; E)$ is k -sparse, then it is k -orientable: the edges of M can be directed so that each vertex has at most k directed edges coming out.

DEF: If M is k -sparse, let

$$X(M) = \{D \subseteq M^2 : (M; D) \text{ is a } k\text{-orientation of } M\} \subseteq \{0, 1\}^{M^2}.$$

Note that this is an $\text{Aut}(M)$ -flow.

Sparse graphs.

DEF: Suppose $k \in \mathbb{N}$. A graph $M = (M; E)$ is k -sparse if for all finite $A \subseteq M$ we have $|E[A]| \leq k|A|$.

FACT: If the graph $M = (M; E)$ is k -sparse, then it is k -orientable: the edges of M can be directed so that each vertex has at most k directed edges coming out.

DEF: If M is k -sparse, let

$$X(M) = \{D \subseteq M^2 : (M; D) \text{ is a } k\text{-orientation of } M\} \subseteq \{0, 1\}^{M^2}.$$

Note that this is an $\text{Aut}(M)$ -flow.

Theorem A

FACT: (Hrushovski) There is an ω -categorical 2-sparse graph M_F with all vertices of infinite valency.

Theorem A' (DE, Jan Hubička and Jaroslav Nešetřil)

Suppose M is a countable, k -sparse graph of infinite valency. If $H \leq \text{Aut}(M)$ is amenable, then H has infinitely many orbits on M^2 .

COROLLARY: There is no precompact amenable subgroup of $\text{Aut}(M_F)$.

Theorem A

FACT: (Hrushovski) There is an ω -categorical 2-sparse graph M_F with all vertices of infinite valency.

Theorem A' (DE, Jan Hubička and Jaroslav Nešetřil)

Suppose M is a countable, k -sparse graph of infinite valency. If $H \leq \text{Aut}(M)$ is amenable, then H has infinitely many orbits on M^2 .

COROLLARY: There is no precompact amenable subgroup of $\text{Aut}(M_F)$.

Theorem A

FACT: (Hrushovski) There is an ω -categorical 2-sparse graph M_F with all vertices of infinite valency.

Theorem A' (DE, Jan Hubička and Jaroslav Nešetřil)

Suppose M is a countable, k -sparse graph of infinite valency. If $H \leq \text{Aut}(M)$ is amenable, then H has infinitely many orbits on M^2 .

COROLLARY: There is no precompact amenable subgroup of $\text{Aut}(M_F)$.

Proof of Thm A': Step 1

- Suppose M is a graph with all vertices of infinite valency and $H \leq \text{Aut}(M)$ has finitely many orbits on M^2 .
- If $c \in M$ let H_c denote the stabilizer of c in H .
- For $c \in M$ let $\text{cl}(c)$ be the union of the finite H_c -orbits on M .
- There is $n \in \mathbb{N}$ s.t. $|\text{cl}(c)| \leq n$ for all $c \in M$.
- If $b \in \text{cl}(c)$ then $\text{cl}(b) \subseteq \text{cl}(c)$.
- STEP 1: There are adjacent $a, b \in M$ such that b is in an infinite H_a -orbit and a is in an infinite H_b -orbit.

PROOF: Suppose there do not exist such a, b . Then for every edge a, b in M either $a \in \text{cl}(b)$ or $b \in \text{cl}(a)$. Take b with $\text{cl}(b)$ of maximal size. There is $a \notin \text{cl}(b)$ adjacent to b . By assumption, $\text{cl}(a) \supset \text{cl}(b)$: contradiction.

Proof of Thm A': Step 1

- Suppose M is a graph with all vertices of infinite valency and $H \leq \text{Aut}(M)$ has finitely many orbits on M^2 .
- If $c \in M$ let H_c denote the stabilizer of c in H .
- For $c \in M$ let $\text{cl}(c)$ be the union of the finite H_c -orbits on M .
- There is $n \in \mathbb{N}$ s.t. $|\text{cl}(c)| \leq n$ for all $c \in M$.
- If $b \in \text{cl}(c)$ then $\text{cl}(b) \subseteq \text{cl}(c)$.
- STEP 1: There are adjacent $a, b \in M$ such that b is in an infinite H_a -orbit and a is in an infinite H_b -orbit.

PROOF: Suppose there do not exist such a, b . Then for every edge a, b in M either $a \in \text{cl}(b)$ or $b \in \text{cl}(a)$. Take b with $\text{cl}(b)$ of maximal size. There is $a \notin \text{cl}(b)$ adjacent to b . By assumption, $\text{cl}(a) \supset \text{cl}(b)$: contradiction.

Proof of Thm A': step 2

- GIVEN: M is a k -sparse graph, $H \leq \text{Aut}(M)$, and $a, b \in M$ are adjacent and such that a is in an infinite H_b -orbit and b is in an infinite H_a -orbit.
- **Show** H is not amenable.
- Suppose there is an H -invariant probability measure μ on $X(M)$.
- Let $S(ab) = \{D \in X(M) : (a, b) \in D\}$. May assume $p = \mu(S(ab)) > 0$.
- Let b_1, \dots, b_n be in the same H_a -orbit as b and s_i the characteristic function of $S(ab_i)$. Note $\mu(S(ab_i)) = p$.
- For $D \in X(M)$,

$$\sum_{i \leq n} s_i(D) \leq k \text{ so } \int_{D \in X(M)} \sum_{i \leq n} s_i(D) d\mu(D) \leq k.$$

- So $np \leq k$: contradiction.

Proof of Thm A': step 2

- GIVEN: M is a k -sparse graph, $H \leq \text{Aut}(M)$, and $a, b \in M$ are adjacent and such that a is in an infinite H_b -orbit and b is in an infinite H_a -orbit.
- **Show** H is not amenable.
- Suppose there is an H -invariant probability measure μ on $X(M)$.
- Let $S(ab) = \{D \in X(M) : (a, b) \in D\}$. May assume $p = \mu(S(ab)) > 0$.
- Let b_1, \dots, b_n be in the same H_a -orbit as b and s_i the characteristic function of $S(ab_i)$. Note $\mu(S(ab_i)) = p$.
- For $D \in X(M)$,

$$\sum_{i \leq n} s_i(D) \leq k \text{ so } \int_{D \in X(M)} \sum_{i \leq n} s_i(D) d\mu(D) \leq k.$$

- So $np \leq k$: contradiction.

Proof of Thm A': step 2

- GIVEN: M is a k -sparse graph, $H \leq \text{Aut}(M)$, and $a, b \in M$ are adjacent and such that a is in an infinite H_b -orbit and b is in an infinite H_a -orbit.
- **Show** H is not amenable.
- Suppose there is an H -invariant probability measure μ on $X(M)$.
- Let $S(ab) = \{D \in X(M) : (a, b) \in D\}$. May assume $p = \mu(S(ab)) > 0$.
- Let b_1, \dots, b_n be in the same H_a -orbit as b and s_i the characteristic function of $S(ab_i)$. Note $\mu(S(ab_i)) = p$.
- For $D \in X(M)$,

$$\sum_{i \leq n} s_i(D) \leq k \text{ so } \int_{D \in X(M)} \sum_{i \leq n} s_i(D) d\mu(D) \leq k.$$

- So $np \leq k$: contradiction.

Proof of Thm A': step 2

- GIVEN: M is a k -sparse graph, $H \leq \text{Aut}(M)$, and $a, b \in M$ are adjacent and such that a is in an infinite H_b -orbit and b is in an infinite H_a -orbit.
- **Show** H is not amenable.
- Suppose there is an H -invariant probability measure μ on $X(M)$.
- Let $S(ab) = \{D \in X(M) : (a, b) \in D\}$. May assume $p = \mu(S(ab)) > 0$.
- Let b_1, \dots, b_n be in the same H_a -orbit as b and s_i the characteristic function of $S(ab_i)$. Note $\mu(S(ab_i)) = p$.
- For $D \in X(M)$,

$$\sum_{i \leq n} s_i(D) \leq k \text{ so } \int_{D \in X(M)} \sum_{i \leq n} s_i(D) d\mu(D) \leq k.$$

- So $np \leq k$: contradiction.

Proof of Thm A': step 2

- GIVEN: M is a k -sparse graph, $H \leq \text{Aut}(M)$, and $a, b \in M$ are adjacent and such that a is in an infinite H_b -orbit and b is in an infinite H_a -orbit.
- **Show** H is not amenable.
- Suppose there is an H -invariant probability measure μ on $X(M)$.
- Let $S(ab) = \{D \in X(M) : (a, b) \in D\}$. May assume $p = \mu(S(ab)) > 0$.
- Let b_1, \dots, b_n be in the same H_a -orbit as b and s_i the characteristic function of $S(ab_i)$. Note $\mu(S(ab_i)) = p$.
- For $D \in X(M)$,

$$\sum_{i \leq n} s_i(D) \leq k \text{ so } \int_{D \in X(M)} \sum_{i \leq n} s_i(D) d\mu(D) \leq k.$$

- So $np \leq k$: contradiction.

Further results

THEOREM B: Suppose $Y \subseteq X(\text{Aut}(M_F))$ is a minimal $\text{Aut}(M_F)$ -subflow. Then all $\text{Aut}(M_F)$ -orbits on Y are meagre in Y .

Other things: Find e.a. subgroups of $\text{Aut}(M_F)$ which are maximal e.a. ; likewise for amenable subgroups...

Further results

THEOREM B: Suppose $Y \subseteq X(\text{Aut}(M_F))$ is a minimal $\text{Aut}(M_F)$ -subflow. Then all $\text{Aut}(M_F)$ -orbits on Y are meagre in Y .

Other things: Find e.a. subgroups of $\text{Aut}(M_F)$ which are maximal e.a. ; likewise for amenable subgroups...

Open Questions

QUESTION: (Bodirsky, . . .) If M is a structure homogeneous for a finite relational language, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

SIDE QUESTION: Is there a homogeneous structure in a finite relational language in which a sparse graph of infinite valency can be interpreted?

QUESTION: (A. Ivanov) If M is ω -categorical and $\text{Aut}(M)$ is amenable, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

Open Questions

QUESTION: (Bodirsky, . . .) If M is a structure homogeneous for a finite relational language, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

SIDE QUESTION: Is there a homogeneous structure in a finite relational language in which a sparse graph of infinite valency can be interpreted?

QUESTION: (A. Ivanov) If M is ω -categorical and $\text{Aut}(M)$ is amenable, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

Open Questions

QUESTION: (Bodirsky, . . .) If M is a structure homogeneous for a finite relational language, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

SIDE QUESTION: Is there a homogeneous structure in a finite relational language in which a sparse graph of infinite valency can be interpreted?

QUESTION: (A. Ivanov) If M is ω -categorical and $\text{Aut}(M)$ is amenable, is there a precompact e.a. subgroup $H \leq \text{Aut}(M)$?

Hrushovski's construction I

- \mathcal{G} : class of finite graphs $(A; R)$
- If $C \subseteq A \in \mathcal{G}$ let

$$\delta(C) = 2|C| - |R[C]|.$$

(Predimension of C .)

- If $A \subseteq B \in \mathcal{C}$ write $A \leq_d B$ if $\delta(X) > \delta(A)$ whenever $A \subset X \subseteq B$.
- Note: If $A \leq_d B \leq_d C$ then $A \leq_d C$.

Hrushovski's construction I

- \mathcal{G} : class of finite graphs $(A; R)$
- If $C \subseteq A \in \mathcal{G}$ let

$$\delta(C) = 2|C| - |R[C]|.$$

(Predimension of C .)

- If $A \subseteq B \in \mathcal{C}$ write $A \leq_d B$ if $\delta(X) > \delta(A)$ whenever $A \subset X \subseteq B$.
- Note: If $A \leq_d B \leq_d C$ then $A \leq_d C$.

Hrushovski's construction I

- \mathcal{G} : class of finite graphs $(A; R)$
- If $C \subseteq A \in \mathcal{G}$ let

$$\delta(C) = 2|C| - |R[C]|.$$

(Predimension of C .)

- If $A \subseteq B \in \mathcal{C}$ write $A \leq_d B$ if $\delta(X) > \delta(A)$ whenever $A \subset X \subseteq B$.
- Note: If $A \leq_d B \leq_d C$ then $A \leq_d C$.

Hrushovski's construction II

- $F : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ an increasing function which tends to infinity.
- Let

$$\mathcal{G}_F = \{A \in \mathcal{G} : \delta(Y) \geq F(|Y|) \text{ for all } Y \subseteq A\}.$$

- For suitable F the class (\mathcal{G}_F, \leq_d) has free amalgamation over \leq_d -substructures.
- In this case the Fraïssé limit construction gives a countable graph M_F characterised by:
 - ▶ M_F is the union of a chain of finite \leq_d -subgraphs;
 - ▶ every graph in \mathcal{G}_F is isomorphic to a \leq_d -subgraph of M_F ;
 - ▶ isomorphisms between finite \leq_d -subgraphs of M_F extend to automorphisms.
- The graph M_F is 2-sparse and ω -categorical.

Hrushovski's construction II

- $F : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ an increasing function which tends to infinity.
- Let

$$\mathcal{G}_F = \{A \in \mathcal{G} : \delta(Y) \geq F(|Y|) \text{ for all } Y \subseteq A\}.$$

- For suitable F the class (\mathcal{G}_F, \leq_d) has free amalgamation over \leq_d -substructures.
- In this case the Fraïssé limit construction gives a countable graph M_F characterised by:
 - ▶ M_F is the union of a chain of finite \leq_d -subgraphs;
 - ▶ every graph in \mathcal{G}_F is isomorphic to a \leq_d -subgraph of M_F ;
 - ▶ isomorphisms between finite \leq_d -subgraphs of M_F extend to automorphisms.
- The graph M_F is 2-sparse and ω -categorical.

Hrushovski's construction II

- $F : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ an increasing function which tends to infinity.
- Let

$$\mathcal{G}_F = \{A \in \mathcal{G} : \delta(Y) \geq F(|Y|) \text{ for all } Y \subseteq A\}.$$

- For suitable F the class (\mathcal{G}_F, \leq_d) has free amalgamation over \leq_d -substructures.
- In this case the Fraïssé limit construction gives a countable graph M_F characterised by:
 - ▶ M_F is the union of a chain of finite \leq_d -subgraphs;
 - ▶ every graph in \mathcal{G}_F is isomorphic to a \leq_d -subgraph of M_F ;
 - ▶ isomorphisms between finite \leq_d -subgraphs of M_F extend to automorphisms.
- The graph M_F is 2-sparse and ω -categorical.